## Regression: Transformations

Grinnell College

October 6, 2025

## Four Assumptions for the Linear Model

- Independence
  - Observations aren't correlated with each other
- Errors are normally distrubted with mean 0
  - Want a random scattering above and below the horizontal line at 0
- Homoskedasticity
  - The general spread of the residuals should be constant through the whole graph
  - ▶ Eg We don't want a megaphone shape pattern in the graph
- X and y's relationship is linear

#### General Plan

Can we do anything with violated assumptions?

- Yes, transformations!
- Logarithms
  - ► (FORMAT: Response Var.'s Scale Explanatory Var.'s Scale)
  - ► Linear-Log
  - Log-Linear
  - ► Log-Log
- Power transformations
  - Square Root (generally of y)
  - Square (generally of x)
- Waaaay more that we can't get into

#### Independence

#### Sometimes data is temporally (or spatially) related

- Eg my chess rating over time
- Have techinques for this
  - Back shift operator: subtract this years ave. September temp from last year's ave. September temp
  - Moving average: our current prediction is the (weighted?) average of the last few time points
  - ► Auto-Regressive: Observations are correlated with their neighbors

#### Sometimes it's phsycially related

- 6 green onions grown in the same small flower pot
- Have techinques for this as well
  - Mixed Models (not in this class)
  - Side step the issue (average over the six green onions)

 Grinnell College
 STA-209
 October 6, 2025
 4 / 35

### Non-Normality

#### Sometimes data just behaves differently

- CHECK FOR LURKING VARIABLE
  - A varible not yet looked at could drive strange behaviro
- I guess 3 heads out of four coin flips; I can only underguess by 1, overguess by 3 (not symmetric!)
  - Logistic Regression
- Discreet data with very few non-0 numbers
  - Discreet distribution over normal, eg Poisson Distribution
  - Zero-Inflated Regression Model
- It's just weird
  - ▶ Bootstrap (later) or simulations (also later)

### Heteroskedasticity

 $\label{lem:common violation of our assumptions is....hetersked a sticity!$ 

### Heteroskedasticity


A common violation of our assumptions is....heterskedasticity!

- It's common the st. dev. of your residuals changes depending on where you are at in the predicted vs residual graph
- Usually (but not always) scientifically expected
- St. Dev. usually (but not always) balloons as the predictions increase


Is there a way to transform the data into a more "usable" form?



#### Example



#### Example: Continued



#### Goal

We want data in a linear form, randomly scattered around a line with constant standard deviation (spread).

- We need a monotonic function that takes in (positive) numeric data
- and makes really large numbers not that large
- while keeping the small numbers relatively small
- and we want to be able to "back transform" (work backwards to get the original data).

Ideas?

 Grinnell College
 STA-209
 October 6, 2025
 11 / 35

#### Goal


We want data in a linear form, randomly scattered around a line with constant standard deviation (spread).

- We need a monotonic function that takes in (positive) numeric data
- and makes really large numbers not that large
- while keeping the small numbers relatively small
- and we want to be able to "back transform" (work backwards to get the original data).


Ideas? The logarithm does this, and so does the (positive) square root

 Grinnell College
 STA-209
 October 6, 2025
 12 / 35

## Example: Continued Some More



## Example: Predictions on Log-Log Scale



### **Back Transforming**


**Back Transforming** is applying a function to an already transformed variable to undo the transformation.

- ▶ Eg taking the square root of a squared variable
- ▶ Eg Undoing the log function by exponentiation
- ▶ Sounds more intimidating than it is

Usually back transformed values do well but not always


- Estimated spread (on the linear scale) balloons as you go up.
- Being off by .1 log units...
  - ightharpoonup when the log value is low it's not bad (eg  $10^{.5}=3.16$  vs  $10^{.6}=3.98$ )
  - when the log value is high is bad (eg  $10^2 = 100$ ,  $10^{2.1} = 126$ )

### Example: Predictions on Linear-Linear Scale



16/35

### Example: Predictions on Linear-Linear Scale



## So what just happened?

Instead of

$$y = \beta_0 + \beta_1 X + e$$

we fit

$$\log(y) = \beta_0 + \beta_1 \log(X) + e$$

Raw Value	Log <sub>10</sub> Value
.2 (=5 <sup>-1</sup> )	698
.5	301
1	0
5	.698
500	2.698

#### What's the catch?

#### Well.....

- We are modeling the log of the response, log(y), and not the response, y.
- Best-fit-line is guaranteed "best" only on the scale modeled
- And we are now fitting the median.....math is weird

Q: Wait, what??


#### Interpretations Background

Well.....we are modeling the log of the response, log(y), and not the response, y.

- We are assumign the errors are bell shaped on the log scale
  - ▶ The distribution will be skewed when we back transform
  - But the median remains the same
- Best-fit-line is guaranteed "best" only on the scale modeled
- And we are now fitting the median....math is weird

Q: Wait, what??

## **Graphic Explanation**



## Estimated Log-Log Regression Equation

$$\log(\hat{y}) = \hat{\beta}_0 + \hat{\beta}_1 * \log(X)$$

$$\hat{y} = \exp(\hat{\beta}_0 + \hat{\beta}_1 * \log(X))$$

$$\hat{y} = \exp(\hat{\beta}_0) * \exp(\hat{\beta}_1 * \log(X))$$

# Log-Log Model's Interpretation for $\hat{eta}_0$

Two Interpretations:

#### **BETTER:**

When the log of the explanatory variable is 0, we expect the median response to be  $\exp(\hat{\beta}_0)$ 

#### WORSE:

When the log of the explanatory variable is 0, we expect the (mean/median) of the log of the response to be  $\hat{\beta}_0$ 

## Slope Interpretation Primer

$$\log(\hat{y}) = \hat{\beta}_0 + \hat{\beta}_1 * \log(X)$$

$$\hat{y} = \exp(\hat{\beta}_0 + \hat{\beta}_1 * \log(X))$$

$$\hat{y} = \exp(\hat{\beta}_0) * \exp(\hat{\beta}_1 * \log(X))$$

$$\hat{y_{new}} = \exp(\hat{\beta_0}) * \exp(\hat{\beta_1} * \log(1.10 * X))$$
 $\hat{y_{new}} = \exp(\hat{\beta_0}) * \exp(\hat{\beta_1} * \log(X)) * \exp(\hat{\beta_1} * \log(1.10))$ 
 $\hat{y_{new}} = \exp(\hat{\beta_0}) * \exp(\hat{\beta_1} * \log(X)) * 1.10^{(\hat{\beta_1})}$ 
 $\hat{y_{new}} = \hat{y} * 1.10^{(\hat{\beta_1})}$ 

# Log-Log Model's Interpretation for $\hat{eta}_1$

#### Two Interpretations

#### BETTER:

When the the explanatory variable increases by 10% we expect the median response to increase by a multiplicative factor of  $1.1^{\hat{\beta}_1}$ 

#### WORSE:

When the log of the explanatory variable increases by 1, we expect the (mean/median) of the log of the response to increase by  $\hat{\beta}_1$ 

#### **Example Continued**

Predicted log(Brain Weight) = 2.13 + .75 \* log(Body Weight)

 $\beta_0$ : For a mammal with a log body weight of 0 (= the body weight of the animal is 1kg), then the predicted (median) weight of it's brain is  $e^2.14 = 8.499g$ 

 $\beta_1$ : If the body weight of the animal increases by 10%, then the predicted (median) brain weight increases by a multiplicative factor of  $1.1^{.75}=1.074$ 

 Grinnell College
 STA-209
 October 6, 2025
 26 / 35

#### **Example Continued**

Artic Fox has a body weight of 3.385kg, what is the predicted median weight for it's brain?

 $Predicted \ log(Brain \ Weight) \ = \ 2.13 \ + \ .75*log(Body \ Weight)$ 

### **Example Continued**

Artic Fox has a body weight of 3.385kg, what is the predicted median weight for it's brain?

$$Predicted log(Brain Weight) = 2.13 + .75 * log(3.385)$$

Predicted 
$$log(Brain Weight) = 3.044$$

Predicted Brain Weight 
$$= e^{(3.044)} = 20.999g$$

Residual = Obs. 
$$-$$
 Predicted =  $44.50 - 20.999 = 23.501$ 

## Log-Linear

Estimate Log-Linear Regression Equation:

$$\log(\hat{y}) = \hat{\beta}_0 + \hat{\beta}_1 * X$$

$$\hat{y} = \exp(\hat{\beta}_0 + \hat{\beta}_1 * X)$$

$$\hat{y} = \exp(\hat{\beta}_0) * \exp(\hat{\beta}_1 * X)$$

 $\hat{\beta_0}$ : Interpretation is the same as for the log log model  $\hat{\beta_0}$ : If the explanatory variable is 0, then the median of the response is exp(), we believe

 $\hat{\beta_1}$ : For a one unit increase in X we expect the median response to increase by a mulitplicative factor of  $\exp(\hat{\beta_1})$ 

 Grinnell College
 STA-209
 October 6, 2025
 29 / 35

### Linear-Log

Estimate Linear-Log Regression Equation:

$$\hat{y} = \hat{\beta_0} + \hat{\beta_1} * \log(X)$$

I don't know of any special interpretations to this one....useful if your x-axis is stretched out waaay too far

 Grinnell College
 STA-209
 October 6, 2025
 30 / 35

## What's the point?


Our goal is to achieve a graph that is linear between whatever is on the x-axis and whatever is on the y-axis with a random scattering points above and below the line (4 assumptions)

- We have techniques help us a way to achieve these goals
- The log() transformation is popular to fix ballooning heteroskedasticity
  - Comes at a cost
  - No "best" properties on the human scale
  - ► The spread of our predictions balloons
  - ▶ Comments can be restricted to medians, not means
- Transformations in general can help us with this goal


#### Other Transformations

- Not uncommon for X to be replaced with X<sup>2</sup>
- The response can be raised to a power
  - Often square root or cubic root
  - ▶ Box-Cox Transformation gives a formula to optimize what power to raise the response to
  - Useful if the units give an indication (see next slide)
- Generalized linear models (GLM's) is a way to deal with the non-normal, quirky data via a different distribution
  - ▶ Eg Logistic and Poisson regression
  - Eg Exponential family
- Finally, non-parametric methods such as ranking the data
  - Calling Spearman's Correlation....

# Algae Blooms



## Algae Blooms: Transformed



#### Next Time

How do we make a linear model when we have categories for our explanatory variable?